

THE MISSISSIPPI KITE

Vol. 36 (1)

July 2006



Mississippi
Ornithological
Society

THE MISSISSIPPI KITE

Application for membership in the Mississippi Ornithological Society may be made to the treasurer. The Mississippi Kite is sent to all members not in arrears for dues. Send change of address, requests for back issues, and claims for undelivered or defective copies to the treasurer.

Information for Authors

The Mississippi Kite publishes original articles that advance the study of birdlife in the state of Mississippi. Submission of articles describing species occurrence and distribution, descriptions of behaviors, notes on the identification of Mississippi birds, as well as scientific studies from all fields of ornithology are encouraged. All manuscripts, in both a hard copy and digital copy format, should be submitted to the editor.

COPY – Paper manuscripts should be *typed, double-spaced* throughout, on high quality paper. Digital copy manuscripts should be prepared using 12 pt. Times New Roman font. If possible, please submit computer files in Microsoft Word although WordPerfect formats will be accepted. Handwritten manuscripts will also be accepted, but please contact the editor prior to submission. Scientific names should be *italicized*.

STYLE – For questions of style consult previous issues of *The Mississippi Kite*. Manuscripts should include a title page (including names and addresses of all authors), text (beginning on page 2), literature cited (if applicable), tables, figure legends (on a separate page), and figures. Number all pages (in the upper right-hand corner) through the tables. Avoid footnotes.

LITERATURE CITED – List all references cited in the text alphabetically by the author's last name in the Literature Cited section. Citations should conform to the style of a recent issue of *The Mississippi Kite*.

NOMENCLATURE - Common names of bird species should be capitalized. Provide the scientific name (*italicized*) at the first mention of each species. Nomenclature should follow the American Ornithologists' Union Checklist of North American Birds, 7th edition (1998) and the 42nd (2000) through 46th (2005) supplements.

TABLES – Tables should be formatted with the size of *The Mississippi Kite* in mind and should be interpretable without reference to the text.

FIGURES – Figures should be appropriate for photoreproduction without retouching.

THE MISSISSIPPI KITE

The Mississippi Kite is a biannual periodical published by the Mississippi Ornithological Society to record and further the study of Mississippi birdlife.

Vol. 36, No. 1

July 2006

CONTENTS

NESTING AND NEST SITE CHARACTERISTICS OF BLACK-CROWNED NIGHT-HERONS (<i>NYCTICORAX NYCTICORAX</i>) AT SWAN LAKE, YAZOO NATIONAL WILDLIFE REFUGE, HOLLANDALE, MISSISSIPPI <i>by Mary P. Stevens and Jerry L. Litton</i>	2
FIRST BREEDING RECORD OF ROSEATE SPOONBILL (<i>PLATALEA AJAJA</i>) IN MISSISSIPPI <i>by Mary P. Stevens and Don McKee</i>	8
MOS OFFICERS AND MEMBERSHIP INFORMATION.....	Back Cover

COVER IMAGE: Mississippi Kite (*Ictinia mississippiensis*) pen and ink drawing by David A. Cimprich.

**NESTING AND NEST SITE CHARACTERISTICS OF
BLACK-CROWNED NIGHT-HERONS (*NYCTICORAX NYCTICORAX*)
AT SWAN LAKE, YAZOO NATIONAL WILDLIFE REFUGE,
HOLLANDALE, MISSISSIPPI**

Mary P. Stevens
5120 Reddoch Drive
Jackson, MS 39211
and
Jerry L. Litton
LightScribe Nature Photography
Pelahatchie, MS 39145

Black-crowned Night-Herons (*Nycticorax nycticorax*) are considered a “species of greatest conservation need” under Mississippi’s Comprehensive Wildlife Conservation Strategy (Mississippi Department of Wildlife, Fisheries and Parks 2005). Black-crowned Night-Herons are widespread and common in North America (Palmer 1962), but very little is known about their nesting habits and habitats in Mississippi (Turcotte and Watts 1999). Black-crowned Night-Herons serve as indicators of environmental quality in wetland habitats because they are high on the food chain, accumulate contaminants, and have a wide geographic distribution (Custer et al. 1991, Davis 1993). Declines in many populations noted in the late 1960s were probably attributable to the use of DDT (Davis 1993).

Black-crowned Night-Herons are known to nest in eight heronries in Mississippi in Holmes, Issaquena, Tallahatchie, Warren, Washington and Yazoo counties (Mississippi Colonial Waterbird Count 1994-2006). Figure 1 is a photograph taken of a Black-crowned Night-Heron at the rookery in Yazoo County. The Washington County colony is located at Swan Lake, Yazoo National Wildlife Refuge, near Hollandale. Swan Lake is a 3,457-acre oxbow lake of the Mississippi River. The heron colony at Swan Lake is situated in a baldcypress-buttonbush swamp and is

protected from human disturbance and habitat destruction, two major causes of the species' declines in the late 1960's and early 1970's (Davis 1993). Since further research is needed to determine the breeding status of this species in Mississippi (Turcotte and Watts 1999), a study of nesting and nest site characteristics of Black-crowned Night-Herons was conducted at the Swan Lake heronry during the spring of 2000. This colony was first censused in 1995 for the Mississippi Colonial Waterbird Count. Twenty-five active Black-crowned Night-Heron nests were found in 1995; 15 nests for each of the years 1996, 1997 and 1998; 50 nests for each of the years 2000, 2001 and 2002; 26 nests for the year 2004; 30 nests for the year 2005; and 50 nests for the year 2006. The colony was not counted in 1999 and 2003. This paper presents additional information regarding nesting activity during the spring of 2000.



Figure 1. Photo taken at the heronry in Yazoo County on May 15, 2006 by Bill Stripling of Vicksburg, Mississippi

METHODS

The study was conducted from March 11, 2000 to May 28, 2000. A total of eleven visits were made to the study site during this time period. Data collection was conducted from a canoe to facilitate quiet travel through the heronry, thereby reducing disturbance to nesting birds (Tremblay and Ellison 1979). An area of 0.85 acres was selected as the study site within a larger mixed-species heronry. Nests were considered active based on the presence of adults at the nest. Each nest monitored was marked with a numbered metal tag. After nests were marked, photographs were taken of nesting birds and their habitat. Each nest tree was identified to species. Tree height was estimated visually. We determined the height of each nest above the water level by establishing a baseline value at one of the nest sites on the first day of the study. All nest heights were measured in inches and keyed to the baseline value. The nest location relative to the tree trunk was recorded, and the nest orientation was taken with a compass. The number of nests per tree and nesting associates for each tree were documented. The number of eggs and young per nest were recorded with the aid of a telescoping mirror used to view the contents of the nest. The number of eggs lost from intact nests by predation or unknown causes was recorded. The date that eggs, incubation and young were first observed was documented.

RESULTS

We located 40 active nests (including some re-nesting attempts), all of which were built in baldcypress. Tree heights ranged from 5-110 feet and most nests were situated slightly over three feet above water level (37.95 inches, range 12-65 inches). Almost all of the nests (38 of 40) were adjacent to the tree trunk, and most that were measured (15 of 25) were facing in a southerly direction (Table 1). All except one of the nests were built in live trees.

Eggs and incubation were first observed on March 25 and young were first seen on April 9. There were usually 3 to 4 nests of Black-crowned Night-Herons and nesting associates per tree. Great Egrets were the most common nesting associate. Of the 33 nest trees surveyed, 31 had Great Egrets nesting in association with Black-crowned Night-Herons, one had Great Egrets, Black-crowned Night-Herons, and Cattle Egrets, and one had only Black-crowned Night-Herons. Seven trees had more than one pair of nesting Black-crowned Night-Herons.

Fifteen of the 40 nests documented in the study area were lost to catastrophic events such as high water or wind. Two nests lost their tags and their disposition could not be determined. The remaining 23 nests lost on average 1.17 eggs per nest due to predation or from uncertain causes. Six of these nests, however, retained complete egg sets throughout incubation and produced a complete set of young. One nest lost the entire clutch (three eggs). These 23 nests hatched an average of 2.35 young per nest (range 0-3).

Table 1. Nest Orientation Determined by Compass

Southeast	Southwest	South	North	Northwest	West	East
3 nests	5 nests	7 nests	2 nests	2 nests	2 nests	4 nests

DISCUSSION

Black-crowned Night-Herons breed on every continent except Australia and Antarctica. As might be expected of a widely distributed species, these birds are flexible in their selection of breeding sites, “[s]o varied as to be difficult to describe – fresh, brackish, and salt-water situations appear equally suitable” (Palmer 1962). Based on the published literature (e.g., Palmer 1962, Jenni 1969, Burger 1978, Wiese 1978, Beaver et al 1980 and Davis 1993), Black-crowned Night-Herons choose locally suitable vegetation for nesting sites. They nest in an enormous

variety of habitat types in mixed species colonies generally over water, and choose nest sites from 2 to 160 feet in height, with relatively small clutch sizes (2-3 eggs). Site selection in Black-crowned Night-Herons may suggest predator avoidance (Davis 1993).

In Mississippi, we found that Black-crowned Night-Herons nest in mixed species colonies over water in secluded wooded swamps, which also suggests that site selection may relate to predator avoidance (Mississippi Colonial Waterbird Count 1995-2006). The colony we studied in Mississippi was a mixed species colony located in a secluded wooded swamp over water. The nests were constructed at heights of 12-65 inches above water and clutch size was relatively small averaging 3 eggs per nest. One nest, however, had a clutch of seven eggs, which is almost double the average clutch size as reported in the literature (Davis 1993). One other clutch size mentioned in the literature also had a clutch of seven eggs (Henny 1972).

LITERATURE CITED

- Beaver, D. L., R. G. Osborn, and T. W. Custer. 1980. Nest-site and colony characteristics of wading birds in selected Atlantic coast colonies. *Wilson Bull.* 92:200-220.
- Burger, J. 1978. The pattern and mechanism of nesting in mixed-species heronries. Pp. 45-58 *In: Wading Birds* (Sprunt et al., Eds). National Audubon Society Research Report No. 7, New York.
- Custer, T. W., B. A. Rattner, H. M. Ohlendorf, and M. J. Melancon. 1991. Herons and egrets as proposed indicators of estuarine contamination in the United States. *Proc. Int. Ornithol. Congr.* 20:2474-2479.

- Davis, W. E. 1993. Black-crowned Night-Heron (*Nycticorax nycticorax*). *In*: The Birds of North America, No. 74 (A. Poole and F. Bill, eds.). Philadelphia: The Academy of Natural Sciences; Washington, D C. : The American Ornithologists' Union.
- Henny, C. J. 1972. An analysis of the population dynamics of selected avian species. U. S. Fish Wildlife Service Research Report No. 1.
- Jenni, D. A. 1969. A study of the ecology of four species of herons during the breeding season at Lake Alice Alachua County, Florida. *Ecol. Monogr.* 39:3.
- Mississippi Colonial Waterbird Count Database. 1995-2006.
- Mississippi Department of Wildlife, Fisheries and Parks. 2005. Mississippi's Comprehensive Wildlife Conservation Strategy 2005-2015. Version 1.1. Jackson, MS: Mississippi Department of Wildlife, Fisheries and Parks, Mississippi Museum of Natural Science, Museum Technical Report No. 117, 2005.
- Palmer, R. S. 1962. Handbook of North American Birds. Vol. I: Loons through Flamingos. Yale University Press.
- Tremblay, J. and L. N. Ellison. 1979. Effects of human disturbance on breeding of Black-crowned Night-Herons. *Auk* 96: 364-369.
- Turcotte, W. H. and Watts, D. 1999. Birds of Mississippi. University Press of Mississippi.
- Wiese, J. H. 1978. Heron nest-site selection and its ecological effects. Pp. 27-34 *In*: Wading Birds (Sprunt et al., Eds.). National Audubon Society Research Report No. 7, New York.

**FIRST BREEDING RECORD OF ROSEATE
SPOONBILL (*PLATALEA AJAJA*) IN MISSISSIPPI**

Mary P. Stevens
5120 Reddoch Drive
Jackson, MS 39211
and
Don McKee
P. O. Box 1445
Dauphin Island, AL 36528

On July 5, 2004, Don McKee and Mary Stevens kayaked into a large multi-species heronry in Yazoo County, locally known as White's Lane, and discovered a Roseate Spoonbill (*Platalea ajaja*) nest with three chicks. This is the first known nesting record of Roseate Spoonbills in Mississippi. The nest was located on the northern border of the heronry. One adult in breeding plumage was present at the nest. The chicks appeared to have primary feather sheaths, so their age was estimated at approximately five days (Dumas 2000). The nest was large and conical and was located in a buttonbush approximately five feet above the water. Photos of the nest were taken on July 5, 2004 by Don McKee and have been deposited with the Mississippi Ornithological Society Records Committee. See Figure 1 below for one of the photos taken on that day. Six subadult spoonbills were observed perched in nearby tupelo trees.

Two subadult spoonbills were observed at this site in June, 2000 by David Linden, biologist at Yazoo National Wildlife Refuge. On June 30, 2004, Mary Stevens, Jenny Thompson, Bryan Fedrick, and Meredith Wilson observed one subadult spoonbill perched in a tupelo tree next to the nest site, but no evidence of nesting was observed at that time. However, considering a 22-day incubation period (Dumas 2000), incubation should have begun around June 8.



Figure 1. Photo of Roseate Spoonbill chick in nest taken on July 5, 2004 by Don McKee at the heronry in Yazoo County, Mississippi

On June 4, 2005, four spoonbills in breeding plumage were observed in the colony and on June 10, 2005, two subadult spoonbills were observed on the south side of the colony. On subsequent visits to the colony during June 2005 and June 2006 to conduct the Mississippi Colonial Waterbird Count, observers found no evidence of nesting of Roseate Spoonbills (Mississippi Colonial Waterbird Count 2000-2006).

This heronry has been selected as one of Mississippi's Important Bird Areas by Audubon Mississippi. The site is a 960-acre baldcypress-tupelo-buttonbush swamp located in Yazoo County at 32° 50' 16"N, 90° 27' 57"W. It is privately owned and

located adjacent to commercial catfish ponds. It is bordered on the west by Panther Swamp National Wildlife Refuge and supports breeding populations of White Ibises, Great Egrets, Snowy Egrets, Little Blue Herons, Cattle Egrets, Black-crowned Night-Herons, Great Blue Herons, Tricolored Herons, Anhingas and Double-crested Cormorants. The Mississippi Colonial Waterbird Count reports approximately 2,000-3,000 nesting pairs of waterbirds annually in this colony. This site has the largest known concentration of nesting White Ibis in the state.

LITERATURE CITED:

Dumas, J. V. 2000. Roseate Spoonbill (*Ajaia ajaja*). In *The Birds of North America*, No. 490 (A. Poole and F. Gill, eds.). The Birds of North America, Inc., Philadelphia, PA.

Mississippi Colonial Waterbird Count Database. White's Lane Colony, Yazoo County, 2000-2006.

THE MISSISSIPPI KITE

(ISSN 0737-0393)

Mississippi Ornithological Society
Organized 30 April 1955

Officers

President	Carl Smith III P. O. Box 227 Stoneville, MS 38776
Vice President	Bill McGehee 114 Pecanwood Dr. Natchez, MS 39120
Secretary	Dianne Patterson 195 Autumn Tr. Columbus, MS 39705
Treasurer	Jan Dubuisson 22410 Glad Acres Pass Christian, MS 39571
Editor: MOS Newsletter	Gene and Shannon Knight 79 Highway 9 West Oxford, MS 38655
Co-Editors: <i>The Mississippi Kite</i>	Margaret Copeland Nick Winstead Marion Schiefer P. O. Box 1505 Starkville, MS 39760

Mississippi Ornithological Society Memberships

Honorary.....	No dues	Student/Senior.....	\$10
Library Subscriptions.....	\$15	Individual.....	\$15
Family.....	\$25	Sustaining.....	\$30
Life.....	\$300		

Membership dues should be sent to the Treasurer.